



MIAMI PEDIATRICS

Visit	Questionnaire	Immunizations	Labs & Screens
Newborn		Hep B*	Transcutaneous bilirubin
Two weeks		Beyfortus*	Transcutaneous bilirubin*
One month	Maternal depression screen		
Two months	Maternal depression screen	Pediarix (DTaP, IPV, Hep B) PCV HiB Rotavirus	
Four months	Maternal depression screen	Pediarix (DTaP, IPV, Hep B) PCV HiB Rotavirus	
Six months	Maternal depression screen	Pediarix (DTaP, IPV, Hep B) PCV HiB Rotavirus Flu** COVID**	
Seven months		Flu** COVID**	
Nine months	Ages & Stages questionnaire		
One year		MMR Hep A	Vision (eye scanner) CBC, lead
Thirteen to Fourteen months		HiB PCV	
Fifteen months		VZV DTaP	
Eighteen months	Ages & Stages questionnaire M-CHAT	Hep A	
Two years	M-CHAT		Vision (eye scanner) CBC*
Thirty months	Ages & Stages questionnaire		
Three years			Vision (eye scanner)
Four years		MMR Kinrix (DTaP, IPV)	Vision (eye scanner) Hearing
Five years		VZV	Vision (eye scanner) Hearing



MIAMI PEDIATRICS

Visit	Questionnaire	Immunizations	Labs & Screens
Six years			Vision (Snellen) Hearing
Seven years			
Eight years			Vision (Snellen) Hearing
Nine years			
Ten years			Lipid profile Vision (Snellen) Hearing
Eleven years		TDaP MCV	CBC (girls)
Twelve years	PHQ-9	HPV	CBC (girls) Vision (Snellen)
Thirteen years	PHQ-9	HPV	CBC (girls) Hearing
Fourteen years	PHQ-9		
Fifteen years	PHQ-9		Vision (Snellen)
Sixteen years	PHQ-9	MCV Men-B	Hearing
Seventeen years	PHQ-9	Men-B	Lipid profile +/- STD testing
Eighteen years	PHQ-9		+/- STD testing^
Nineteen years	PHQ-9		+/- STD testing
Twenty years	PHQ-9		+/- STD testing Hearing
Twenty-one years	PHQ-9	TDaP	+/- STD testing

*Only if needed

**Influenza and COVID vaccines are optional annually starting at 6 months of age

^After age 18 years, if drawing blood for STD testing, include Hepatitis C at least 1 time